	SBAK Communication Iool	SBAR Communication Iool
S	Situation: This is, and I am the nurse for patienton pediatrics. I am concerned about the patient because(give concise description — Pain, VS change, status change, increasing PEWS score).	Situation: This is, and I am the nurse for patienton pediatrics. I am concerned about the patient because(give concise description — Pain, VS change, status change, increasing PEWS score).
B	Background: Patient was admitted onwith Recent procedures include (OR, tests, investigations) The child's condition has changed in the lastminutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS	Background: Patient was admitted onwith Recent procedures include (OR, tests, investigations) The child's condition has changed in the lastminutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS
A	Assessment: I think the problem is I have given (O2, analgesia, stopped infusion) The patient seems to be unstable, deteriorating, needs intervention.	Assessment: I think the problem is I have given (O2, analgesia, stopped infusion)  The patient seems to be unstable, deteriorating, needs intervention.
R	Recommendation: I need you to:  • Come and see the patient now  • Call the MRP or pediatrician  • Order tests or labs (CXR, Labs)  Is there anything I need to do now (monitoring, fluids)  End call by confirming plan of care	Recommendation: I need you to:  • Come and see the patient now  • Call the MRP or pediatrician  • Order tests or labs (CXR, Labs)  Is there anything I need to do now (monitoring, fluids)  End call by confirming plan of care
	SBAR Communication Tool	SBAR Communication Tool
S	Situation: This is, and I am the nurse for patient on pediatrics. I am concerned about the patient because (give concise description — Pain, VS change, status change, increasing PEWS score).	Situation: This is, and I am the nurse for patienton pediatrics. I am concerned about the patient because (give concise description — Pain, VS change, status change, increasing PEWS score).
B	Background: Patient was admitted onwith Recent procedures include (OR, tests, investigations) The child's condition has changed in the lastminutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS	Background: Patient was admitted onwith Recent procedures include (OR, tests, investigations) The child's condition has changed in the lastminutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS
A	Assessment: I think the problem is I have given (O2, analgesia, stopped infusion) The patient seems to be unstable, deteriorating, needs intervention.	Assessment: I think the problem is I have given (O2, analgesia, stopped infusion) The patient seems to be unstable, deteriorating, needs intervention.
R	Recommendation: I need you to:  • Come and see the patient now  • Call the MRP or pediatrician  • Order tests or labs (CXR, Labs)  Is there anything I need to do now (monitoring, fluids)  End call by confirming plan of care	Recommendation: I need you to:  Come and see the patient now Call the MRP or pediatrician Order tests or labs (CXR, Labs) Is there anything I need to do now (monitoring, fluids) End call by confirming plan of care
	SBAR Communication Tool	SBAR Communication Tool
S	Situation: This is, and I am the nurse for patienton pediatrics. I am concerned about the patient because (give concise	Situation: This is, and I am the nurse for patienton pediatrics. I am concerned about the patient because (give concise
	description — Pain, VS change, status change, increasing PEWS score).	description — Pain, VS change, status change, increasing PEWS score).
В	· ————————————————————————————————————	
B A	description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted on with Recent procedures include (OR, tests, investigations) The child's condition has changed in the last minutes. The latest vital	description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted on with Recent procedures include (OR, tests, investigations) The child's condition has changed in the last minutes. The latest vital
B A R	description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted on with Recent procedures include (OR, tests, investigations) The child's condition has changed in the last minutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS  Assessment: I think the problem is I have given (O2, analgesia, stopped infusion)	description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted on with Recent procedures include (OR, tests, investigations) The child's condition has changed in the last minutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS  Assessment: I think the problem is I have given (O2, analgesia, stopped infusion)
A	description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted on with Recent procedures include (OR, tests, investigations) The child's condition has changed in the last minutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS  Assessment: I think the problem is I have given (O2, analgesia, stopped infusion)  The patient seems to be unstable, deteriorating, needs intervention.  Recommendation: I need you to:  Come and see the patient now  Call the MRP or pediatrician	description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted onwith Recent procedures include (OR, tests, investigations) The child's condition has changed in the lastminutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS  Assessment: I think the problem is I have given (O2, analgesia, stopped infusion) The patient seems to be unstable, deteriorating, needs intervention.  Recommendation: I need you to: Come and see the patient now Call the MRP or pediatrician
A	description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted on with Recent procedures include (OR, tests, investigations) The child's condition has changed in the last minutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS  Assessment: I think the problem is I have given (O2, analgesia, stopped infusion)  The patient seems to be unstable, deteriorating, needs intervention.  Recommendation: I need you to:  Come and see the patient now  Call the MRP or pediatrician  Order tests or labs (CXR, Labs)  Is there anything I need to do now (monitoring, fluids)  End call by confirming plan of care	description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted on with Recent procedures include (OR, tests, investigations) The child's condition has changed in the last minutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS  Assessment: I think the problem is I have given (O2, analgesia, stopped infusion)  The patient seems to be unstable, deteriorating, needs intervention.  Recommendation: I need you to:  • Come and see the patient now  • Call the MRP or pediatrician  • Order tests or labs (CXR, Labs)  Is there anything I need to do now (monitoring, fluids)  End call by confirming plan of care
A	description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted on with Recent procedures include (OR, tests, investigations) The child's condition has changed in the last minutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS  Assessment: I think the problem is I have given (O2, analgesia, stopped infusion)  The patient seems to be unstable, deteriorating, needs intervention.  Recommendation: I need you to:  • Come and see the patient now  • Call the MRP or pediatrician  • Order tests or labs (CXR, Labs)  Is there anything I need to do now (monitoring, fluids)	description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted onwith Recent procedures include (OR, tests, investigations) The child's condition has changed in the lastminutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS  Assessment: I think the problem is I have given (O2, analgesia, stopped infusion) The patient seems to be unstable, deteriorating, needs intervention.  Recommendation: I need you to: Come and see the patient now Call the MRP or pediatrician Order tests or labs (CXR, Labs) Is there anything I need to do now (monitoring, fluids)
A	description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted on with Recent procedures include (OR, tests, investigations) The child's condition has changed in the last minutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS  Assessment: I think the problem is I have given (O2, analgesia, stopped infusion)  The patient seems to be unstable, deteriorating, needs intervention.  Recommendation: I need you to:  • Come and see the patient now  • Call the MRP or pediatrician  • Order tests or labs (CXR, Labs)  Is there anything I need to do now (monitoring, fluids)  End call by confirming plan of care  SBAR Communication Tool  Situation: This is, and I am the nurse for patient on pediatrics. I am concerned about the patient because (give concise)	description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted onwith Recent procedures include (OR, tests, investigations) The child's condition has changed in the lastminutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS  Assessment: I think the problem is I have given (O2, analgesia, stopped infusion) The patient seems to be unstable, deteriorating, needs intervention.  Recommendation: I need you to:
A R	description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted on with Recent procedures include (OR, tests, investigations) The child's condition has changed in the last minutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS  Assessment: I think the problem is I have given (O2, analgesia, stopped infusion)  The patient seems to be unstable, deteriorating, needs intervention.  Recommendation: I need you to:  • Come and see the patient now  • Call the MRP or pediatrician  • Order tests or labs (CXR, Labs)  Is there anything I need to do now (monitoring, fluids)  End call by confirming plan of care  SBAR Communication Tool  Situation: This is, and I am the nurse for patient on pediatrics. I am concerned about the patient because (give concise description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted on with Recent procedures include (OR, tests, investigations) The child's condition has changed in the last minutes. The latest vital	description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted onwith Recent procedures include (OR, tests, investigations) The child's condition has changed in the lastminutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS  Assessment: I think the problem is I have given (O2, analgesia, stopped infusion) The patient seems to be unstable, deteriorating, needs intervention.  Recommendation: I need you to: Come and see the patient now Call the MRP or pediatrician Order tests or labs (CXR, Labs) Is there anything I need to do now (monitoring, fluids) End call by confirming plan of care  SBAR Communication Tool  Situation: This is, and I am the nurse for patient on pediatrics. I am concerned about the patient because (give concise description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted on with Recent procedures include (OR, tests, investigations) The child's condition has changed in the last minutes. The latest vital
A R	description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted on with Recent procedures include (OR, tests, investigations) The child's condition has changed in the last minutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS  Assessment: I think the problem is I have given (O2, analgesia, stopped infusion)  The patient seems to be unstable, deteriorating, needs intervention.  Recommendation: I need you to:  • Come and see the patient now  • Call the MRP or pediatrician  • Order tests or labs (CXR, Labs)  Is there anything I need to do now (monitoring, fluids)  End call by confirming plan of care  SBAR Communication Tool  Situation: This is, and I am the nurse for patient on pediatrics. I am concerned about the patient because (give concise description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted on with Recent procedures include (OR, tests, investigations) The child's condition has changed in the last minutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS  Assessment: I think the problem is I have given (O2, analgesia, stopped infusion)	description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted onwith Recent procedures include (OR, tests, investigations) The child's condition has changed in the lastminutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS  Assessment: I think the problem is I have given (O2, analgesia, stopped infusion) The patient seems to be unstable, deteriorating, needs intervention.  Recommendation: I need you to:  • Come and see the patient now  • Call the MRP or pediatrician  • Order tests or labs (CXR, Labs)  Is there anything I need to do now (monitoring, fluids)  End call by confirming plan of care  SBAR Communication Tool  Situation: This is, and I am the nurse for patient on pediatrics. I am concerned about the patient because (give concise description — Pain, VS change, status change, increasing PEWS score).  Background: Patient was admitted on with Recent procedures include (OR, tests, investigations) The child's condition has changed in the last minutes. The latest vital signs are HR, RR, BP, O2 sats, Temp, Pain Score, PEWS  Assessment: I think the problem is I have given (O2, analgesia, stopped infusion)

### **Brighton Pediatric Early Warning Score**

	0	1	2	3	
Behavior	Playing Appropriate	Sleeping	Irritable	Lethargic &/OR Confused &/OR Reduced response to pain	
Respiratory	Within normal parameters, no recession OR tracheal tug	10 above normal parameters, using accessory muscles &/OR 30+% FiO2 or 4+ liters/min	>20 above normal parameters, recessing/retractions, tracheal tug &/OR 40+ % FiO2 or 6+ liters/min	5 below normal parameters with sterna recessions/retractions, tracheal tug or grunting &/OR 50% FiO2 or 8+ liters/min	
Cardiovascular	Pink/normal &/OR Capillary refill 1–2 seconds	Pale &/OR Capillary refill 3 seconds	Grey <b>&amp;/OR</b> Capillary refill 4 seconds, Tachycardia of 20 above normal rate	Grey & mottled or capillary refill 5 seconds or above <b>OR</b> Tachycardia of 30 above normal rate of bradycardia	
	Bronchodilators e	very 20 minutes <b>&amp;/OR</b> Pe	ersistent vomiting follow	ing surgery (2 points each)	
	TOTAL PEWS SCORE:				

PHSA330 | APR.2019

# **Brighton Pediatric Early Warning Score**

	0	1	2	3	
Behavior	Playing Appropriate	Sleeping	Irritable	Lethargic &/OR Confused &/OR Reduced response to pain	
Respiratory	Within normal parameters, no recession OR tracheal tug	10 above normal parameters, using accessory muscles &/OR 30+% FiO2 or 4+ liters/min	>20 above normal parameters, recessing/retractions, tracheal tug &/OR 40+ % FiO2 or 6+ liters/min	5 below normal parameters with sterna recessions/retractions, tracheal tug or grunting &/OR 50% FiO2 or 8+ liters/min	
Cardiovascular	Pink/normal &/OR Capillary refill 1–2 seconds	Pale &/OR Capillary refill 3 seconds	Grey <b>&amp;/OR</b> Capillary refill 4 seconds, Tachycardia of 20 above normal rate	Grey & mottled or capillary refill 5 seconds or above <b>OR</b> Tachycardia of 30 above normal rate of bradycardia	
	Bronchodilators e	very 20 minutes <b>&amp;/OR</b> Pe	ersistent vomiting follow	ing surgery (2 points each)	
	TOTAL PEWS SCORE:				

PHSA330 | APR.2019

#### **Brighton Pediatric Early Warning Score**

	0	1	2	3		
Behavior	Playing Appropriate	Sleeping	Irritable	Lethargic &/OR Confused &/OR Reduced response to pain		
Respiratory	Within normal parameters, no recession <b>OR</b> tracheal tug	10 above normal parameters, using accessory muscles &/OR 30+% FiO2 or 4+ liters/min	>20 above normal parameters, recessing/retractions, tracheal tug &/OR 40+ % FiO2 or 6+ liters/min	5 below normal parameters with sterna recessions/retractions, tracheal tug or grunting &/OR 50% FiO2 or 8+ liters/min		
Cardiovascular	Pink/normal &/OR Capillary refill 1–2 seconds	Pale &/OR Capillary refill 3 seconds	Grey &/OR Capillary refill 4 seconds, Tachycardia of 20 above normal rate	Grey & mottled or capillary refill 5 seconds or above <b>OR</b> Tachycardia of 30 above normal rate of bradycardia		
	Bronchodilators e	very 20 minutes <b>&amp;/OR</b> Pe	rsistent vomiting follow	ing surgery (2 points each)		
	TOTAL DEWIS SCORE.					

PHSA330 | APR.2019

# **Brighton Pediatric Early Warning Score**

brighten reductie Early Warning Score				
	0	1	2	3
Behavior	Playing Appropriate	Sleeping	Irritable	Lethargic &/OR Confused &/OR Reduced response to pain
Respiratory	Within normal parameters, no recession <b>OR</b> tracheal tug	10 above normal parameters, using accessory muscles &/OR 30+ % FiO2 or 4+ liters/min	>20 above normal parameters, recessing/retractions, tracheal tug &/OR 40+ % FiO2 or 6+ liters/min	5 below normal parameters with sterna recessions/retractions, tracheal tug or grunting &/OR 50% FiO2 or 8+ liters/min
Cardiovascular	Pink/normal &/OR Capillary refill 1–2 seconds	Pale <b>&amp;/OR</b> Capillary refill 3 seconds	Grey &/OR Capillary refill 4 seconds, Tachycardia of 20 above normal rate	Grey & mottled or capillary refill 5 seconds or above <b>OR</b> Tachycardia of 30 above normal rate of bradycardia
	Bronchodilators	every 20 minutes <b>&amp;/OR</b> Pe	ersistent vomiting follow	ing surgery (2 points each)
TOTAL PEWS SCORE:				

**Brighton Pediatric Early Warning Score** 

	0	1	2	3	
Behavior	Playing Appropriate	Sleeping	Irritable	Lethargic <b>&amp;/OR</b> Confused <b>&amp;/OR</b> Reduced response to pain	
Respiratory	Within normal parameters, no recession <b>OR</b> tracheal tug	10 above normal parameters, using accessory muscles &/OR 30+% FiO2 or 4+ liters/min	>20 above normal parameters, recessing/retractions, tracheal tug &/OR 40+ % FiO2 or 6+ liters/min	5 below normal parameters with sterna recessions/retractions, tracheal tug or grunting &/OR 50% FiO2 or 8+ liters/min	
Cardiovascular	Pink/normal &/OR Capillary refill 1–2 seconds	Pale <b>&amp;/OR</b> Capillary refill 3 seconds	Grey <b>&amp;/OR</b> Capillary refill 4 seconds, Tachycardia of 20 above normal rate	Grey & mottled or capillary refill 5 seconds or above <b>OR</b> Tachycardia of 30 above normal rate of bradycardia	
	Bronchodilators e	very 20 minutes <b>&amp;/OR</b> Pe	rsistent vomiting follow	ing surgery (2 points each)	
TOTAL PEWS SCORE:					

PHSA330 | APR.2019

### **Brighton Pediatric Early Warning Score**

	0	1	2	3		
Behavior	Playing Appropriate	Sleeping	Irritable	Lethargic &/OR Confused &/OR Reduced response to pain		
Respiratory	Within normal parameters, no recession <b>OR</b> tracheal tug	10 above normal parameters, using accessory muscles &/OR 30+ % FiO2 or 4+ liters/min	>20 above normal parameters, recessing/retractions, tracheal tug &/OR 40+ % FiO2 or 6+ liters/min	5 below normal parameters with sterna recessions/retractions, tracheal tug or grunting &/OR 50% FiO2 or 8+ liters/min		
Cardiovascular	Pink/normal &/OR Capillary refill 1–2 seconds	Pale <b>&amp;/OR</b> Capillary refill 3 seconds	Grey &/OR Capillary refill 4 seconds, Tachycardia of 20 above normal rate	Grey & mottled or capillary refill 5 seconds or above <b>OR</b> Tachycardia of 30 above normal rate of bradycardia		
	Bronchodilators e	every 20 minutes &/OR Pe	ersistent vomiting follow	ing surgery (2 points each)		
	TOTAL PEWS SCORE:					

PHSA330 | APR.2019

#### **Brighton Pediatric Early Warning Score**

	0	1	2	3	
Behavior	Playing Appropriate	Sleeping	Irritable	Lethargic &/OR Confused &/OR Reduced response to pain	
Respiratory	Within normal parameters, no recession <b>OR</b> tracheal tug	10 above normal parameters, using accessory muscles &/OR 30+% FiO2 or 4+ liters/min	>20 above normal parameters, recessing/retractions, tracheal tug &/OR 40+ % FiO2 or 6+ liters/min	5 below normal parameters with sterna recessions/retractions, tracheal tug or grunting &/OR 50% FiO2 or 8+ liters/min	
Cardiovascular	Pink/normal <b>&amp;/OR</b> Capillary refill 1–2 seconds	Pale <b>&amp;/OR</b> Capillary refill 3 seconds	Grey <b>&amp;/OR</b> Capillary refill 4 seconds, Tachycardia of 20 above normal rate	Grey & mottled or capillary refill 5 seconds or above <b>OR</b> Tachycardia of 30 above normal rate of bradycardia	
	Bronchodilators e	very 20 minutes <b>&amp;/OR</b> Pe	rsistent vomiting follow	ing surgery (2 points each)	
	TOTAL PEWS SCORE:				

PHSA330 | APR.2019

# **Brighton Pediatric Early Warning Score**

	0	1	2	3
Behavior	Playing Appropriate	Sleeping	Irritable	Lethargic &/OR Confused &/OR Reduced response to pain
Respiratory	Within normal parameters, no recession <b>OR</b> tracheal tug	10 above normal parameters, using accessory muscles &/OR 30+% FiO2 or 4+ liters/min	>20 above normal parameters, recessing/retractions, tracheal tug &/OR 40+ % FiO2 or 6+ liters/min	5 below normal parameters with sterna recessions/retractions, tracheal tug or grunting &/OR 50% FiO2 or 8+ liters/min
Cardiovascular	Pink/normal <b>&amp;/OR</b> Capillary refill 1–2 seconds	Pale <b>&amp;/OR</b> Capillary refill 3 seconds	Grey <b>&amp;/OR</b> Capillary refill 4 seconds, Tachycardia of 20 above normal rate	Grey & mottled or capillary refill 5 seconds or above <b>OR</b> Tachycardia of 30 above normal rate of bradycardia
	Bronchodilators e	very 20 minutes <b>&amp;/OR</b> Pe	ersistent vomiting follow	ing surgery (2 points each)
				TOTAL PEWS SCORE:

PHSA330 | APR.2019 PHSA330 | APR.2019